## Properties of Ionic Salts in Supercritical CHF3-CH3OH Mixtures

D.L. Goldfarb<sup>1,2,</sup>, D.P. Fernández<sup>2</sup> and H.R. Corti<sup>1</sup>

<sup>1</sup>Unidad Actividad Química CNEA Av. del Libertador 8250 1429 Buenos Aires, Argentina

<sup>2</sup>INQUIMAE Universidad de Buenos Aires Ciudad Universitaria Pab. II 1428 Buenos Aires, Argentina

The dielectric constant and the molar conductivity of solutions of the ionic salt tetrabutylammonium tetraphenylborate (TBATPB) in supercritical mixtures of CHF<sub>3</sub>-CH<sub>3</sub>OH were measured with a high-pressure coaxial capacitor at 308 K and 323 K, and pressures up to 15 MPa. The formation of a second phase was explored in the capacitor and compared with independent measurements. The solubility of CH<sub>3</sub>OH in CHF<sub>3</sub> and the solubility of TBATPB in CHF<sub>3</sub>-CH<sub>3</sub>OH mixtures were measured by UV spectroscopy or constant pressure sampling followed by chemical analysis. The density of the mixtures was measured by a vibrating tube densimeter as a function of composition.

The dielectric constant of the mixture was analyzed using different theories, while the conductivity was described in terms of ionic mobilities and ionic speciation.